



EMBRACE 3.6L

For Use on Wheat to suppress Fusarium head blight (scab)

SECTION 18 SPECIFIC EXEMPTIONS

File Symbol: 07MT04

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF MONTANA UNDER SECTION 18 SPECIFIC EXEMPTIONS

These use directions must be in possession of the user at the time of pesticide application. It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Read entire Directions for Use and Disclaimer of Warranties before using this product. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions including precautions when spraying in the vicinity of aquatic areas, on the registered product label for EMBRACE 3.6L (EPA Reg No. 1381-203).

Any adverse effects resulting from the use of EMBRACE 3.6L under this specific exemption must be immediately reported to the State of Montana Department of Agriculture.

EFFECTIVE PERIOD:

These use directions becomes effective 5/14/07 and expires on 7/20/07.

DIRECTIONS FOR USE

A maximum of 300,500 acres of Wheat may be treated under this exemption.

RECOMMENDED APPLICATIONS

CROP	DISEASES	RATE OF EMBRACE 3.6L
Wheat (including durum)	<i>Fusarium</i> head blight (scab)	4 fl oz per acre

USE DIRECTIONS

Apply EMBRACE 3.6L as a preventive foliar spray. For optimum disease suppression, the lowest rate of a spray surfactant should be tank-mixed with EMBRACE 3.6L. EMBRACE 3.6L may be applied in a minimum of 10 gallons of spray solution per acre by ground sprayer or in a minimum of 3 gallons of spray solution per acre by aircraft spray equipment.

Application through any type of irrigation equipment is prohibited.

Do not make more than 1 application of a tebuconazole containing product to wheat per season. One application may be made using ground or aerial equipment at a rate of 4 fluid ounces of formulated product (1.8 oz a.i.) per acre per year. EMBRACE 3.6L may be applied up through full head emergence (Feekes 10.51). Application may not be made within 30 days of harvest. Straw cut after harvest may be fed or used for bedding. Do not enter treated field for 12 hours after application.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to freshwater, estuarine and marine fish and invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark. Runoff may be hazardous to aquatic organisms in neighboring areas. Do not contaminate water when disposing of equipment wash water or rinsate.

For emergencies call CHEMTREC toll free 24 Hours A Day 1-800-424-9300

For PRODUCT USE Information visit at www.agrilance.com

As with any crop protection product, always read and follow label directions.

Distributed By:
Agrilance, LLC
P.O. Box 64089
St. Paul, MN 55164-0089